



Escherichia coli de infección urinaria de origen comunitario

Total	N°		Sexo	Edad	Antibiótico	Porcentaje	
	N°	Intermedia				Intermedia	Resistente
125	2	1	M	≤14 años	AMK	1,60	0,80
129	0	39			GEN	0,00	30,23
128	1	98			AMP	0,78	76,56
39	5	14			AMC	12,82	35,90
111	15	60			SAM	13,51	54,05
109	20	56			CEP	18,35	51,38
115	1	33			CXM	0,87	28,70
136	0	90			SXT	0,00	66,18
122	1	40			CIP	0,82	32,79
136	6	11			NIT	4,41	8,09
932	12	13	F	≤14 años	AMK	1,29	1,39
957	9	166			GEN	0,94	17,35
951	9	720			AMP	0,95	75,71
386	64	107			AMC	16,58	27,72
899	182	340			SAM	20,24	37,82
730	173	264			CEP	23,70	36,16
214	0	46			CZO	0,00	21,50
890	9	186			CXM	1,01	20,90
1013	0	604			SXT	0,00	59,62
889	3	2,14			CIP	0,34	0,24
1049	49	51			NIT	4,67	4,86




CENTRO DE REFERENCIA NACIONAL DE RESISTENCIA A LOS ANTIMICROBIANOS

Total Nº	Nº		Sexo	Edad	Antibiótico	Porcentaje	
	Intermedia	Resistente				Intermedia	Resistente
354	6	13	M	15 a 60	AMK	1,69	3,67
370	6	131			GEN	1,62	35,41
327	6	285			AMP	1,83	87,16
148	33	70			AMC	22,30	47,30
348	68	184			SAM	19,54	52,87
247	39	154			CEP	15,79	62,35
189	1	45			CZO	1,12	50,56
299	16	137			CXM	5,35	45,82
349	0	247			SXT	0,00	70,77
365	3	247			CIP	0,82	67,67
386	14	63			NIT	3,63	16,32
4033	81	52	F	15 a 60	AMK	2,01	1,29
4105	82	755			GEN	2,00	18,39
3794	61	2671			AMP	1,61	70,40
2288	410	618			AMC	17,92	27,01
3470	645	1201			SAM	18,59	34,61
2715	581	977			CEP	21,40	35,99
1203	2	16			CZO	0,17	1,33
3564	89	866			CXM	2,50	24,30
4222	30	2229			SXT	0,71	52,79
4084	33	1862			CIP	0,81	45,59
4527	258	344			NIT	5,70	7,60



CENTRO DE REFERENCIA NACIONAL DE RESISTENCIA A LOS ANTIMICROBIANOS


Total	N°		Sexo	Edad	Antibiótico	Porcentaje	
	N°	Intermedia				Intermedia	Resistente
572	16	15	M	> 60	AMK	2,80	2,62
610	12	243			GEN	1,97	39,84
555	8	448			AMP	1,44	80,72
232	41	112			AMC	17,67	48,28
554	88	297			SAM	15,88	53,61
473	81	285			CEP	17,12	60,25
111	0	59			CZO	0,00	53,15
520	35	246			CXM	6,73	47,31
615	3	406			SXT	0,49	66,02
566	5	448			CIP	0,88	79,15
656	42	149			NIT	6,40	22,71
2230	38	29	F	> 60	AMK	1,70	1,30
2363	47	508			GEN	1,99	21,50
2193	32	1579			AMP	1,46	72,00
1065	189	304			AMC	17,75	28,54
2185	437	739			SAM	20,00	33,82
1563	306	642			CEP	19,58	41,07
681	1	208			CZO	0,15	30,54
2109	80	618			CXM	3,79	29,30
2408	10	1428			SXT	0,42	59,30
2319	21	1350			CIP	0,91	58,21
2561	120	241			NIT	4,69	9,41



CENTRO DE REFERENCIA NACIONAL DE RESISTENCIA A LOS ANTIMICROBIANOS

Escherichia coli de origen hospitalario


Total	N°		Antibiótico	Porcentaje	
	N°	Intermedia	Resistente	Intermedia	Resistente
7341	139	154	AMK	1,89	2,10
6996	84	2120	GEN	1,20	30,30
4542	82	3670	AMP	1,81	80,80
2265	487	888	AMC	21,50	39,21
4504	374	419	TZP	8,30	9,30
5055	20	86	IMP	0,40	1,70
6315	19	95	MEM	0,30	1,50
3124	578	1690	CEP	18,50	54,10
997	0	398	CZO	0,00	39,92
3928	55	1669	CTX	1,40	42,49
7012	7	3254	CAZ	0,10	46,41
6748	47	3158	FEP	0,70	46,80
3974	60	1896	FOX	1,51	47,71
1095	74	658	NAL	6,76	60,09
7469	60	4250	CIP	0,80	56,90
5004	25	3183	SXT	0,50	63,61
4628	324	407	NIT	7,00	8,79
3662	66	352	FOS	1,80	9,61
1845	0	24	COL	0,00	1,30
3048	34	3	TGC	1,12	0,10



CT y TGC solo por CIM

Staphylococcus spp. de origen comunitario

Total	Nº		Especie	Antibiótico	Porcentaje	
Nº	Intermedio	Resistente			Intermedio	Resistente
703	23	61	S. aureus	GEN	3,27	8,68
664	13	196		OXA	1,96	29,52
282	0	98		FOX	0,00	34,75
742	7	237		SXT	0,94	31,94
699	30	67		CIP	4,29	9,59
42	1	1		TEC	2,38	2,38
514	0	0		VAN	0,00	0,00
770	32	87		CLI	4,16	11,30
767	72	314		ERI	9,39	40,94
543	1	27		RIF	0,18	4,97
635	0	3		LNZ	0,00	0,47
58	3	1		MNO	5,17	1,72
593	4	121		TCY	0,67	20,40
637	66	33	Staphylococcus coagulasa negativa	GEN	10,36	5,18
583	2	381		OXA	0,34	65,35
242	0	141		FOX	0,00	58,26
664	2	376		SXT	0,30	56,63
606	44	202		CIP	7,26	33,33
43	1	9		TEC	2,33	20,93
485	0	0		VAN	0,00	0,00
602	39	236		CLI	6,48	39,20
601	18	461		ERI	3,00	76,71
494	5	49		RIF	1,01	9,92
557	0	14		LNZ	0,00	2,51
96	2	6		MNO	2,08	6,25
506	7	142		TCY	1,38	28,06




*VAN solo por CIM

CENTRO DE REFERENCIA NACIONAL DE RESISTENCIA A LOS ANTIMICROBIANOS

Staphylococcus spp. de origen hospitalario

Total	Nº		Especie	Antibiótico	Porcentaje	
Nº	Intermedia	Resistente			Intermedia	Resistente
2196	105	163	S. aureus	GEN	4,78	7,42
2025	4	804		OXA	0,20	39,70
574	0	262		FOX	0,00	45,64
2172	13	1066		SXT	0,60	49,08
2019	50	204		CIP	2,48	10,10
195	1	8		TEC	0,51	4,10
1877	0	0		VAN	0,00	0,00
2231	38	319		CLI	1,70	14,30
2248	121	879		ERI	5,38	39,10
1887	8	96		RIF	0,42	5,09
2119	0	13		LNZ	0,00	0,61
250	13	7		MNO	5,20	2,80
1732	10	414		TCY	0,58	23,90
3269	356	1330	Staphylococcus coagulasa negativa	GEN	10,89	40,69
3032	6	2420		OXA	0,20	79,82
930	0	766		FOX	0,00	82,37
3078	15	2232		SXT	0,49	72,51
2832	195	1427		CIP	6,89	50,39
457	7	22		TEC	1,53	4,81
2774	0	0		VAN	0,00	0,00
3371	101	1827		CLI	3,00	54,20
3384	68	2853		ERI	2,01	84,31
2806	25	519		RIF	0,89	18,50
3246	0	58		LNZ	0,00	1,79
757	7	8		MNO	0,92	1,06
2369	14	609		TCY	0,59	25,71

*VAN solo por CIM




CENTRO DE REFERENCIA NACIONAL DE RESISTENCIA A LOS ANTIMICROBIANOS

Enterobacter cloacae de origen hospitalario

Total	N°		Antibiótico	Porcentaje	
N°	Intermedia	Resistente		Intermedia	Resistente
846	5	18	AMK	0,59	2,13
835	7	198	GEN	0,84	23,71
744	67	117	TZP	9,01	15,73
802	19	24	IPM	2,37	2,99
835	3	24	MEM	0,36	2,87
850	17	278	CAZ	2,00	32,71
116	7	55	CTX	6,03	47,41
799	12	308	CRO	1,50	38,55
843	53	188	FEP	6,29	22,30
243	3	92	SXT	1,23	37,86
32	8	12	NAL	25,00	37,50
846	54	100	CIP	6,38	11,82
96	6	13	FOS	6,25	13,54
470	0	17	COL	0,00	3,62
695	68	40	TGC	9,78	5,76


CT y TGC solo por CIM



CENTRO DE REFERENCIA NACIONAL DE RESISTENCIA A LOS ANTIMICROBIANOS

Klebsiella pneumoniae de origen hospitalario


Total	N°		Antibiótico	Porcentaje	
	N°	Intermedia	Resistente	Intermedia	Resistente
4082	66	442	AMK	1,62	10,83
4052	55	1916	GEN	1,36	47,29
829	171	448	AMC	20,63	54,04
3616	116	2638	AMS	3,21	72,95
3401	279	1505	TZP	8,20	44,25
3632	27	1132	IPM	0,74	31,17
3921	22	1279	MEM	0,56	32,62
831	13	626	CEP	1,56	75,33
4021	13	2822	CAZ	0,32	70,18
1072	15	737	CTX	1,40	68,75
3853	8	2685	CRO	0,21	69,69
3987	24	2805	FEP	0,60	70,35
3121	11	2235	FOX	0,35	71,61
326	43	169	NAL	13,19	51,84
4144	595	1843	CIP	14,36	44,47
1004	275	496	NIT	27,39	49,40
890	86	177	FOS	9,66	19,89
2231	0	187	COL	0,00	8,38
2907	658	348	TGC	22,64	11,97



CT y TGC solo por CIM

Enterococcus spp. de origen hospitalario

Total	N°		Microorganismo	Antibiótico	Porcentaje	
N°	Intermedia	Resistente			Intermedia	Resistente
403	1	50	<i>Enterococcus faecalis</i>	GEH	0,25	12,41
302	0	45		STH	0,00	14,90
805	0	48		AMP	0,00	5,96
759	11	19		LNZ	1,45	2,50
66	0	1		TEC	0,00	1,52
778	9	9		VAN	1,16	1,16
89	0	21	<i>Enterococcus faecium</i>	GEH	0,00	23,60
75	0	18		STH	0,00	24,00
205	0	176		AMP	0,00	85,85
202	0	4		LNZ	0,00	1,98
17	0	1		TEC	0,00	1/17*
203	2	36		VAN	0,99	17,73



* La resistencia se presenta en fracción debido al bajo número de aislamientos analizados.


	Resistencia Natural
--	---------------------



Acinetobacter baumannii complex de origen hospitalario

Total Nº	Nº		Antibiótico	Porcentaje	
	Intermedia	Resistente		Intermedia	Resistente
182	3	77	AMK	1,65	42,31
483	37	199	GEN	7,66	41,20
393	51	190	SAM	12,98	48,35
429	19	292	TZP	4,43	68,07
458	4	268	IPM	0,87	58,52
468	8	279	MEM	1,71	59,62
478	86	246	CAZ	17,99	51,46
481	30	300	FEP	6,24	62,37
126	1	91	SXT	0,79	72,22
483	4	312	CIP	0,83	64,60
280	0	1	COL	0,00	0,36
406	87	41	TGC	21,43	10,10

CT y TGC solo por CIM



Pseudomonas aeruginosa de origen hospitalario

Total	N°		Antibiótico	Porcentaje	
N°	Intermedia	Resistente		Intermedia	Resistente
1644	68	162	AMK	4,14	9,85
1635	131	401	GEN	8,01	24,53
1454	175	395	TZP	12,04	27,17
1556	21	530	IPM	1,35	34,06
1562	79	463	MEM	5,06	29,64
250	40	55	ATM	16,00	22,00
1661	180	352	CAZ	10,84	21,19
1661	148	364	FEP	8,91	21,91
912	0	26	COL	0,00	2,85
1658	103	554	CIP	6,21	33,41

CT y TGC solo por CIM

